



Certain Subclasses of Bi-Univalent and Meromorphic Functions Defined By Al-Oboudi Differential Operator

D. D. Bobalade ¹ and N. D. Sangle ²

¹Department of Mathematics, Shivaji University, Kolhapur, Maharashtra, India- 416004

²Department of Mathematics, D.Y. Patil College of Engineering and Technology
 Kasaba Bawada, Kolhapur, Maharashtra, India 416006

Email: ¹dnyaneshwar.bobalade@gmail.com, ²navneet_sangle@rediffmail.com

Abstract: In this paper, we introduce a subclass $\Sigma_M^{h,p}(k, \delta, \mu, \lambda)$ of bi-univalent and meromorphic functions by using Al-Oboudi differential operator on $\Delta = \{z \in \mathbb{C}: 1 < |z| < \infty\}$. Also we obtain bounds of coefficients $|b_0|$ and $|b_1|$ for functions belongs to $\Sigma_M^{h,p}(k, \delta, \mu, \lambda)$. The results obtained in this paper are more better and generalized of previous results of various author.

Keywords: Harmonic functions, Univalent functions, Differential operator, Extreme points.

2010 Mathematics Subject Classification: 30C45, 30C50.

1. Introduction

Let Σ be the class of functions f of the form

$$f(z) = z + \sum_{l=2}^{\infty} \frac{b_l}{z^l}, \quad (1)$$

which are meromorphic univalent in the domain $\Delta = \{z \in \mathbb{C}: 1 < |z| < \infty\}$. Since every function f belong to Σ has an inverse function f^{-1} exist and inverse function satisfies conditions:

$$f^{-1}(f(z)) = z \quad (z \in \Delta)$$

and

$$f(f^{-1}(w)) = w, \quad w \in \Delta \quad (M < |w| < \infty, M > 0),$$

where

$$f^{-1}(w) = q(w) = w - b_0 - \frac{b_1}{w} - \frac{b_2 + b_0 b_1}{w^2} - \frac{b_3 + 2b_0 b_2 + b_0^2 b_1 + b_1^2}{w^3} + \dots \quad (2)$$

A function $f \in \Sigma$ is said to be meromorphic bi-univalent in Δ if both f and f^{-1} are meromorphic univalent in Δ . The class of meromorphic bi-univalent functions of the form (1) in Δ is denoted by Σ_M .

Srivastava et al. [17], Safa Salehian and Ahmad Zireh [12], Hamidi et al. [7], Amol Patil and Uday Naik [11] and many other researchers (see [4, 5, 8, 9, 10, 14, 15, 18]) have introduced new subclasses of meromorphically bi-univalent functions and obtained estimates on the initial coefficients for functions in each of these subclasses.

Let \mathcal{A} denote the class of analytic functions $h(z)$ of the form

$$h(z) = z + \sum_{l=2}^{\infty} a_l z^l \quad (3)$$

defined in the unit disc $\mathbb{U} = \{z \in \mathbb{C}: |z| < 1\}$ with normalization $h(0) = h_z(0) - 1 = 0$. Let the class of all normalized analytic univalent functions in the unit disc \mathbb{U} is denoted by S . A function $h \in \mathcal{A}$ is said to

be bi-univalent in \mathbb{U} if both h and h^{-1} are univalent in \mathbb{U} . Let the class of analytic bi-univalent functions is denoted by Σ' . Brannan and Taha [2], Srivastava et al. [16] and many other researchers (see [3, 6]) introduced certain subclasses of bi-univalent function class Σ' .

Now, Al-Oboudi [1] introduced the Al-Oboudi operator $D_\delta^k: \mathcal{A} \rightarrow \mathcal{A}$ and defined as

$$D^k h(z) = D_\delta^k h(z) = z + \sum_{l=2}^{\infty} [1 + (l-1)\delta]^k a_l z^l, \quad k \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \delta \geq 0,$$

where $h \in \mathcal{A}$ of the form (3).

Amol Patil and Uday Naik [11] extend the Al-Oboudi operator $D_\delta^k: \Sigma \rightarrow \Sigma$ and defined as

$$D^k f(z) = D_\delta^k f(z) = z + (1-\delta)^k b_0 + \sum_{l=1}^{\infty} [1 - (l+1)\delta]^k b_l z^{-l}, \quad k \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \delta > 1,$$

where $f \in \Sigma$ of the form (1).

In 2019, Saideh Hajiparvaneh and Ahmad Zireh [13] define the subclass $\Sigma_M^{h,p}(\mu, \lambda)$ consisting of meromorphic functions $f(z)$ of the form (1) satisfies the following conditions:

$$f \in \Sigma_M, \left[(1-\lambda) \left(\frac{f(z)}{z} \right)^\mu + \lambda f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} \right] \in h(\Delta)$$

and

$$\left[(1-\lambda) \left(\frac{q(w)}{w} \right)^\mu + \lambda q'(w) \left(\frac{q(w)}{w} \right)^{\mu-1} \right] \in p(\Delta),$$

where q is function given by (2).

Motivated by the aforecited works, we introduce new subclasses of bi-univalent and meromorphic functions by using Al-Oboudi Differential operator. Also obtain the coefficient bounds $|b_0|$ and $|b_1|$ for functions in this new subclasses.

2. Coefficient Estimates

Definition 2.1 Let the analytic functions $h, p: \Delta \rightarrow \mathbb{C}$ be

$$h(z) = 1 + \frac{h_1}{z} + \frac{h_2}{z^2} + \frac{h_3}{z^3} + \dots, \quad p(z) = 1 + \frac{p_1}{z} + \frac{p_2}{z^2} + \frac{p_3}{z^3} + \dots,$$

such that

$$\min\{\Re(h(z)), \Re(p(z))\} > 0 \quad (z \in \Delta).$$

Definition 2.2 A function $f(z)$ of the form (1) is said to be in the class $\Sigma_M^{h,p}(k, \delta, \mu, \lambda)$, $k \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \lambda \geq 1, \lambda > \mu, \mu \geq 0$ and $\delta > 1$ if satisfies the following conditions :

$$f \in \Sigma_M, \left[(1-\lambda) \left(\frac{D_\delta^k f(z)}{z} \right)^\mu + \lambda (D_\delta^k f(z))' \left(\frac{D_\delta^k f(z)}{z} \right)^{\mu-1} \right] \in h(\Delta) \tag{4}$$

and

$$\left[(1-\lambda) \left(\frac{D_\delta^k q(w)}{w} \right)^\mu + \lambda (D_\delta^k q(w))' \left(\frac{D_\delta^k q(w)}{w} \right)^{\mu-1} \right] \in p(\Delta), \tag{5}$$

where q is the function given by (2).

For $k = 0$, the class $\Sigma_M^{h,p}(k, \delta, \mu, \lambda)$ become $\Sigma_M^{h,p}(\mu, \lambda)$, studied by Saideh Hajiparvaneh and Ahmad Zireh [13].

Remark 2.1 For various choices of h and p , we get various subclasses of class $\Sigma_M^{h,p}(k, \delta, \mu, \lambda)$ as follows:

If take $h(z) = p(z) = \left(\frac{1+\frac{1}{z}}{1-\frac{1}{z}} \right)^\alpha = 1 + \frac{2\alpha}{z} + \frac{2\alpha^2}{z^2} + \dots$ ($0 < \alpha \leq 1, z \in \Delta$) in Definition 2.2, then

we get subclass $\Sigma_M^{h,p}(k, \delta, \mu, \lambda) = \Sigma_M^*(k, \delta, \mu, \lambda, \alpha)$, studied by Bobalade and Sangle [4].

Definition 2.3 [4] A function $f(z) \in \Sigma_M^{h,p}(k, \delta, \mu, \lambda)$ of the form (1) belongs to the class $\Sigma_M^*(k, \delta, \mu, \lambda, \alpha)$ if

$$f \in \Sigma_M, \left| \arg \left[(1-\lambda) \left(\frac{D_\delta^k f(z)}{z} \right)^\mu + \lambda (D_\delta^k f(z))' \left(\frac{D_\delta^k f(z)}{z} \right)^{\mu-1} \right] \right| < \frac{\alpha\pi}{2} \quad (z \in \Delta)$$

and

$$\left| \arg \left[(1 - \lambda) \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu} + \lambda (D_{\delta}^k q(w))' \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu-1} \right] \right| < \frac{\alpha\pi}{2} \quad (w \in \Delta),$$

where $k \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \lambda \geq 1, \lambda > \mu, \mu \geq 0, \delta > 1$ and $0 < \alpha \leq 1$.

Remark 2.2 If take $h(z) = p(z) = \frac{1 + \frac{(1-2\beta)z}{1-z}}{1-\frac{1}{z}} = 1 + \frac{2(1-\beta)}{z} + \frac{2(1-\beta)}{z^2} + \dots$ ($0 \leq \beta < 1, z \in \Delta$) in Definition

2.2, then we get subclass $\Sigma_M^{h,p}(k, \delta, \mu, \lambda) = \Sigma_M^*(k, \delta, \mu, \lambda, \beta)$, studied by Bobalade and Sangle [4].

Definition 2.4 [4] A function $f(z) \in \Sigma_M^{h,p}(k, \delta, \mu, \lambda)$ of the form (1) belongs to the class $\Sigma_M^*(k, \delta, \mu, \lambda, \beta)$ if

$$f \in \Sigma_M, \Re \left[(1 - \lambda) \left(\frac{D_{\delta}^k f(z)}{z} \right)^{\mu} + \lambda (D_{\delta}^k f(z))' \left(\frac{D_{\delta}^k f(z)}{z} \right)^{\mu-1} \right] > \beta \quad (z \in \Delta)$$

and

$$\Re \left[(1 - \lambda) \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu} + \lambda (D_{\delta}^k q(w))' \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu-1} \right] > \beta \quad (w \in \Delta),$$

where $k \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \lambda \geq 1, \lambda > \mu, \mu \geq 0, \delta > 1$ and $0 \leq \beta < 1$.

If we put $k = 0$ in the classes $\Sigma_M^*(k, \delta, \mu, \lambda, \alpha)$ and $\Sigma_M^*(k, \delta, \mu, \lambda, \beta)$, then we get two classes $\Sigma_M^*(\mu, \lambda, \alpha)$ and $\Sigma_M^*(\mu, \lambda, \beta)$ respectively, studied by Orhan et al. [10].

Theorem 2.1 Let $f(z)$ of the form (1) belong to the class $\Sigma_M^{h,p}(k, \delta, \mu, \lambda)$. Then

$$|b_0| \leq \min \left[\sqrt{\frac{|h_1|^2 + |p_1|^2}{2(\mu - \lambda)^2(1 - \delta)^{2k}}}, \sqrt{\frac{|h_2| + |p_2|}{|(\mu - 2\lambda)(\mu - 1)(1 - \delta)^{2k}|}} \right] \quad (6)$$

and

$$|b_1| \leq \min \left[\frac{|h_2| + |p_2|}{2|(\mu - 2\lambda)(1 - 2\delta)^k|}, \sqrt{\frac{|h_2|^2 + |p_2|^2}{2(\mu - 2\lambda)^2(1 - 2\delta)^{2k}} + \frac{(\mu - 1)^2(|h_1|^2 + |p_1|^2)^2}{16(\mu - \lambda)^4(1 - 2\delta)^{2k}}} \right]. \quad (7)$$

Proof. From conditions (4) and (5), we have

$$(1 - \lambda) \left(\frac{D_{\delta}^k f(z)}{z} \right)^{\mu} + \lambda (D_{\delta}^k f(z))' \left(\frac{D_{\delta}^k f(z)}{z} \right)^{\mu-1} = h(z) \quad (z \in \Delta) \quad (8)$$

and

$$(1 - \lambda) \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu} + \lambda (D_{\delta}^k q(w))' \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu-1} = p(w) \quad (w \in \Delta), \quad (9)$$

where $h(z)$ and $p(w)$ are functions such that it's real part positive in Δ and have forms

$$h(z) = 1 + \frac{h_1}{z} + \frac{h_2}{z^2} + \frac{h_3}{z^3} + \dots \quad (10)$$

and

$$p(w) = 1 + \frac{p_1}{w} + \frac{p_2}{w^2} + \frac{p_3}{w^3} + \dots \quad (11)$$

Implies

$$(1 - \lambda) \left(\frac{D_{\delta}^k f(z)}{z} \right)^{\mu} + \lambda (D_{\delta}^k f(z))' \left(\frac{D_{\delta}^k f(z)}{z} \right)^{\mu-1} = 1 + \frac{h_1}{z} + \frac{h_2}{z^2} + \frac{h_3}{z^3} + \dots, \quad (z \in \Delta) \quad (12)$$

and

$$(1 - \lambda) \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu} + \lambda (D_{\delta}^k q(w))' \left(\frac{D_{\delta}^k q(w)}{w} \right)^{\mu-1} = 1 + \frac{p_1}{w} + \frac{p_2}{w^2} + \frac{p_3}{w^3} + \dots \quad (w \in \Delta). \quad (13)$$

Now, equating the coefficients in equation (12) and (13), we obtain

$$(\mu - \lambda)(1 - \delta)^k b_0 = h_1, \quad (14)$$

$$(\mu - 2\lambda) \left[(1 - 2\delta)^k b_1 + \left(\frac{\mu-1}{2} \right) (1 - \delta)^{2k} b_0^2 \right] = h_2, \quad (15)$$

$$-(\mu - \lambda)(1 - \delta)^k b_0 = p_1 \tag{16}$$

and

$$(\mu - 2\lambda) \left[-(1 - 2\delta)^k b_1 + \left(\frac{\mu-1}{2}\right) (1 - \delta)^{2k} b_0^2 \right] = p_2. \tag{17}$$

From equation (14) and equation (17), we get

$$h_1 = -p_1 \tag{18}$$

and

$$2(\mu - \lambda)^2 (1 - \delta)^{2k} b_0^2 = h_1^2 + p_1^2. \tag{19}$$

By adding equation (15) to equation (17), we get

$$(\mu - 2\lambda)(\mu - 1)(1 - \delta)^{2k} b_0^2 = h_2 + p_2. \tag{20}$$

Therefore, From equation (19), we get

$$b_0^2 = \frac{h_1^2 + p_1^2}{2(\mu - \lambda)^2 (1 - \delta)^{2k}} \tag{21}$$

and from equation (20), we get

$$b_0^2 = \frac{h_2 + p_2}{(\mu - 2\lambda)(\mu - 1)(1 - \delta)^{2k}}. \tag{22}$$

Hence, from (21) and (22), we find that

$$|b_0|^2 \leq \frac{|h_1|^2 + |p_1|^2}{2(\mu - \lambda)^2 (1 - \delta)^{2k}}$$

and

$$|b_0|^2 \leq \frac{|h_2| + |p_2|}{|(\mu - 2\lambda)(\mu - 1)(1 - \delta)^{2k}|}.$$

Hence

$$|b_0| \leq \min \left[\sqrt{\frac{|h_1|^2 + |p_1|^2}{2(\mu - \lambda)^2 (1 - \delta)^{2k}}}, \sqrt{\frac{|h_2| + |p_2|}{|(\mu - 2\lambda)(\mu - 1)(1 - \delta)^{2k}|}} \right].$$

Now, subtracting equation (17) from equation (15), we obtain

$$2(\mu - 2\lambda)(1 - 2\delta)^k b_1 = h_2 - p_2. \tag{23}$$

By squaring and adding equations (15) and (17), we get new equation. Using equation (19) in new equation, we obtain

$$b_1^2 = \frac{h_2^2 + p_2^2}{2(\mu - 2\lambda)^2 (1 - 2\delta)^{2k}} + \frac{(\mu - 1)^2 (h_1^2 + p_1^2)^2}{16(\mu - \lambda)^4 (1 - 2\delta)^{2k}}. \tag{24}$$

By using equations (10), (11) in equation (23) and (24), finally we yield

$$|b_1| \leq \frac{|h_2| + |p_2|}{2|(\mu - 2\lambda)(1 - 2\delta)^k|} \tag{25}$$

and

$$|b_1| \leq \sqrt{\frac{|h_2|^2 + |p_2|^2}{2(\mu - 2\lambda)^2 (1 - 2\delta)^{2k}} + \frac{(\mu - 1)^2 (|h_1|^2 + |p_1|^2)^2}{16(\mu - \lambda)^4 (1 - 2\delta)^{2k}}}. \tag{26}$$

Hence

$$|b_1| \leq \min \left[\frac{|h_2| + |p_2|}{2|(\mu - 2\lambda)(1 - 2\delta)^k|}, \sqrt{\frac{|h_2|^2 + |p_2|^2}{2(\mu - 2\lambda)^2 (1 - 2\delta)^{2k}} + \frac{(\mu - 1)^2 (|h_1|^2 + |p_1|^2)^2}{16(\mu - \lambda)^4 (1 - 2\delta)^{2k}}} \right]. \tag{27}$$

This complete the proof.

3. Corollaries and Consequences

If we take $k = 0$ in Theorem 2.1, then obtain following Corollary.

Corollary 3.1 [13] *Let $f(z)$ of the form (1) be in the class $\Sigma_M^{h,p}(\mu, \lambda)$, $\lambda \geq 1, \lambda > \mu, \geq 0$. Then*

$$|b_0| \leq \min \left[\sqrt{\frac{|h_1|^2 + |p_1|^2}{2(\mu - \lambda)^2}}, \sqrt{\frac{|h_2| + |p_2|}{|(\mu - 2\lambda)(\mu - 1)|}} \right]$$

and
$$|b_1| \leq \min \left[\frac{|h_2|+|p_2|}{2|(\mu-2\lambda)|}, \sqrt{\frac{|h_2|^2+|p_2|^2}{2(\mu-2\lambda)^2} + \frac{(\mu-1)^2(|h_1|^2+|p_1|^2)^2}{16(\mu-\lambda)^4}} \right].$$

Remark 3.1 Corollary 3.1 is an improvement result of result obtained by Orhan [10] in Theorem 1.2.

If we take $h(z) = p(z) = \left(\frac{1+\frac{1}{z}}{1-\frac{1}{z}}\right)^\alpha = 1 + \frac{2\alpha}{z} + \frac{2\alpha^2}{z^2} + \dots$ ($0 < \alpha \leq 1, z \in \Delta$), in Theorem 2.1, then we obtain following result.

Corollary 3.2 Let $f(z)$ of the form (1) belong to the class $\Sigma_M^*(k, \delta, \mu, \lambda, \alpha)$, then

$$|b_0| \leq \min \left[\frac{2\alpha}{|(\mu-\lambda)(1-\delta)^k|}, \frac{2\alpha}{\sqrt{|(\mu-2\lambda)(\mu-1)(1-\delta)^{2k}|}} \right]$$

and

$$|b_1| \leq \min \left[\frac{2\alpha^2}{|(\mu-2\lambda)(1-2\delta)^k|}, 2\alpha^2 \sqrt{\frac{1}{(\mu-2\lambda)^2(1-2\delta)^{2k}} + \frac{(\mu-1)^2}{(\mu-\lambda)^4(1-2\delta)^{2k}}} \right].$$

Remark 3.2 Corollary 3.2 is an improvement result of result obtained by Bobalade and Sangle [4] in Theorem 2.6.

If we take $k = 0$ in Corollary 3.2, then we obtain following Corollary.

Corollary 3.3 [13] Let $f(z)$ of the form (1) belong to the class $\Sigma_M^*(\mu, \lambda, \beta)$, then

$$|b_0| \leq \min \left[\frac{2\alpha}{(\lambda-\mu)}, \frac{2\alpha}{\sqrt{|(\mu-2\lambda)(\mu-1)|}} \right]$$

and

$$|b_1| \leq \min \left[\frac{2\alpha^2}{(2\lambda-\mu)}, 2\alpha^2 \sqrt{\frac{1}{(\mu-2\lambda)^2} + \frac{(\mu-1)^2}{(\mu-\lambda)^4}} \right].$$

If we take $h(z) = p(z) = \frac{1+\frac{(1-2\beta)}{z}}{1-\frac{1}{z}} = 1 + \frac{2(1-\beta)}{z} + \frac{2(1-\beta)^2}{z^2} + \dots$ ($0 \leq \beta < 1, z \in \Delta$), in Theorem 2.1, then we obtain following result.

Corollary 3.4 Let $f(z)$ of the form (1) be in the class $\Sigma_M^*(k, \delta, \mu, \lambda, \beta)$, then

$$|b_0| \leq \min \left[\frac{2(1-\beta)}{|(\mu-\lambda)(1-\delta)^k|}, \sqrt{\frac{4(1-\beta)}{|(\mu-2\lambda)(\mu-1)(1-\delta)^{2k}|}} \right]$$

and

$$|b_1| \leq \min \left[\frac{2(1-\beta)}{|(\mu-2\lambda)(1-2\delta)^k|}, 2(1-\beta) \sqrt{\frac{1}{(\mu-2\lambda)^2(1-2\delta)^{2k}} + \frac{(1-\beta)^2(\mu-1)^2}{(\mu-\lambda)^4(1-2\delta)^{2k}}} \right].$$

Remark 3.3 Corollary 3.4 is an improvement result of result obtained by Bobalade and Sangle [4] in Theorem 2.3.

If we take $k = 0$ in Corollary 3.4, then we get following Corollary .

Corollary 3.5 [13] Let $f(z)$ of the form (1) belong to the class $\Sigma_M^*(\mu, \lambda, \beta)$, then

$$|b_0| \leq \min \left[\frac{2(1-\beta)}{(\lambda-\mu)}, 2 \sqrt{\frac{(1-\beta)}{|(\mu-2\lambda)(\mu-1)|}} \right]$$

and

$$|b_1| \leq \min \left[\frac{2(1-\beta)}{(2\lambda-\mu)}, 2(1-\beta) \sqrt{\frac{1}{(\mu-2\lambda)^2} + \frac{(1-\beta)^2(\mu-1)^2}{(\mu-\lambda)^4}} \right].$$

Acknowledgment:

The authors wish to thank the referee, for the careful reading of the paper and for the helpful suggestions and comments.

References

- [1] Ai-Oboudi, F. M. (2004). On univalent functions defined by a generalized Salagean operator. *Int.J. of Math. And Math. Sci.*, **27**: 1429-1436.
- [2] Brannan, D. A. and Taha, T. S. (1986). On some classes of bi-univalent functions. *Studia Univ. Babeş-Bolyai Math.*, **31**(2): 70-77.
- [3] Bobalade, D. D. and Sangle, N. D. (2019). Coefficient estimates for a certain subclass of bi-univalent functions defined by multiplier transformation. *International Journal of Research and Analytical Reviews*, **6**(1): 123-128.
- [4] Bobalade, D. D. and Sangle, N. D. (2021). Certain subclasses of bi-univalent and meromorphic functions associated with al-oboudi differential operator. *International Journal of Mathematics Trends and Technology*, **67**(5): 86-89.
- [5] Duren, P. L. (1983). Univalent functions (Grundlehren der Mathematischen Wissenschaften, Band 259, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo).
- [6] Frasin, B. A. and Aouf, M. K. (2011). New subclasses of bi-univalent functions. *Appl. Math. Lett.*, **24**: 1569-1573.
- [7] Hamidi, S. G., Halim, S. A. and Jahangiri, J. M. (2013). Coefficient estimates for a class of meromorphic bi-univalent functions. *C. R. Acad. Sci. Paris, Ser. I*, **351**: 349-352.
- [8] Janani, T. and Murugusundaramoorthy, G. (2014). Coefficient estimates of meromorphic bi-starlike functions of complex order. *Internat. J. Anal. Appl.*, **4**(1): 68-77.
- [9] Kubota, Y. (1977). Coefficients of meromorphic univalent functions. *Kodai Mathematical Seminar Report*, **28**: 253-261.
- [10] Orhan, H., Mangesh, N. and Balaji, V. K. (2014). Initial coefficient bounds for certain subclasses of meromorphic bi-univalent functions. *Asian European J.Math.*, **7**(1): 1-9.
- [11] Patil, A.B. and Naik, U.H. (2017). Coefficient estimates for a new subclass of meromorphic bi-univalent functions defined by Al-Oboudi differential operator. *Global Journal of Pure and Applied Mathematics*, **13**(9): 4405-4414.
- [12] Safa S. and Ahmad Z. (2017). Coefficient estimates for certain subclass of meromorphic and bi-univalent functions. *Commun. Korean Math. Soc.*, **32**(2): 389-397.
- [13] Saideh H. and Ahmad Z. (2019). Coefficients estimates for certain subclass of meromorphic and bi-univalent functions. *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, **68**(1): 654-662.
- [14] Schiffer, M. (1938). Sur un probleme d'extremum de la representation conforme. *Bull. Soc. Math. France*, **66**: 48-55.
- [15] Springer, G. (1951). The Coefficients problem for Schlicht mappings of the exterior of the unit circle. *Trans. Amer. Math. Soc.*, **70**: 421-450.
- [16] Srivastava, H. M., Mishra, A. K. and Gochhayat, P. (2010). Certain subclasses of analytic and bi-univalent functions. *Applied Mathematics Letters*, **23**: 1188-1192.
- [17] Srivastava, H. M., Joshi, S.B., Joshi, S. S. and Haridas, P. (2016). Coefficient Estimates for Certain Subclasses of Meromorphically Bi-Univalent Functions. *Palestine Journal of Mathematics*, **5**(1): 250-258.
- [18] Suzeini, A.H., Samaneh, G. H. and Ravichandran, V. (2011). Coefficient Estimates for Meromorphic Bi-Univalent Functions. *arXiv:1108.4089v1*, 1-9.

□□